

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07C 249/02		A3	(11) International Publication Number: WO 99/03803
			(43) International Publication Date: 28 January 1999 (28.01.99)
(21) International Application Number: PCT/US98/14253		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 8 July 1998 (08.07.98)		Published <i>With international search report.</i>	
(30) Priority Data: 08/892,278 14 July 1997 (14.07.97) US			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/892,278 (CON) Filed on 14 July 1997 (14.07.97)		(88) Date of publication of the international search report: 8 April 1999 (08.04.99)	
(71) Applicant (for all designated States except US): CATALYTICA PHARMACEUTICALS, INC. [US/US]; 430 Ferguson Drive, Mountain View, CA 94043-5272 (US).			
(72) Inventor; and (75) Inventor/Applicant (for US only): GRUBER, John, M. [US/US]; 505 Cypress Point Drive, Mountain View, CA 94043 (US).			
(74) Agents: CIOTTI, Thomas, E. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).			
(54) Title: PROCESS FOR PREPARING AMINOARYLACETYLENES			
(57) Abstract <p>This invention provides a process for preparing aminoarylacetylenes comprising reacting an N-arylmethylidene aminoarylhalide with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt to produce a novel N-arylmethylidene aminoarylacetylene, and hydrolyzing the N-arylmethylidene aminoarylacetylene to the aminoarylacetylene. In one embodiment, the invention provides a process for preparing aminophenylacetylenes comprising reacting an N-benzylidene aminophenylhalide with a terminal acetylene in the presence of a base and the catalyst system to produce a novel N-benzylidene aminophenylacetylene, and hydrolyzing the N-benzylidene aminophenylacetylene to the aminophenylacetylene.</p>			

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07C	A2	(11) International Publication Number: WO 99/03803 (43) International Publication Date: 28 January 1999 (28.01.99)
(21) International Application Number: PCT/US98/14253 (22) International Filing Date: 8 July 1998 (08.07.98) (30) Priority Data: 08/892,278 14 July 1997 (14.07.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/892,278 (CON) Filed on 14 July 1997 (14.07.97) (71) Applicant (for all designated States except US): CATALYTICA PHARMACEUTICALS, INC. [US/US]; 430 Ferguson Drive, Mountain View, CA 94043-5272 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): GRUBER, John, M. [US/US]; 505 Cypress Point Drive, Mountain View, CA 94043 (US). (74) Agents: CIOTTI, Thomas, E. et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PROCESS FOR PREPARING AMINOARYLACETYLENES (57) Abstract This invention provides a process for preparing aminoarylacetylenes comprising reacting an N-arylmethylidene aminoarylhalide with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt to produce a novel N-arylmethylidene aminoarylacetylene, and hydrolyzing the N-arylmethylidene aminoarylacetylene to the aminoarylacetylene. In one embodiment, the invention provides a process for preparing aminophenylacetylenes comprising reacting an N-benzylidene aminophenylhalide with a terminal acetylene in the presence of a base and the catalyst system to produce a novel N-benzylidene aminophenylacetylene, and hydrolyzing the N-benzylidene aminophenylacetylene to the aminophenylacetylene.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROCESS FOR PREPARING AMINOARYLACETYLENES

FIELD OF THE INVENTION

5

This invention relates generally to preparing aminoarylacetylenes. More specifically, it relates to preparing aminoarylacetylenes from aminoarylhalides and terminal acetylenes and to novel compounds which are useful in their preparation. Aminophenylacetylenes are valuable as precursors to pharmaceutically active compounds and to acetylene terminated
10 polymers. In particular, 3-aminophenylacetylene is used as an end-capping agent for high performance polyimide resins.

BACKGROUND OF THE INVENTION

15

Sabourin, *Prepr. Div. Pet. Chem., Am. Chem. Soc.*, vol. 24, pp. 233-239 discloses the preparation of 2-methyl-4-(3-aminophenyl)-3-butyne-2-ol (an aminophenylacetylene) and 3-aminophenylacetylene in two and three steps, respectively, from 3-bromonitrobenzene and 2-methyl-3-butyne-2-ol. In the first step, 3-bromonitrobenzene and 2-methyl-3-butyne-2-ol were reacted in the presence of a catalyst system of bis(triphenylphosphine)palladium
20 dichloride, additional triphenylphosphine, and cuprous iodide in triethylamine solvent at the reflux temperature to obtain 2-methyl-4-(3-nitrophenyl)-3-butyne-2-ol. In the second step, this nitrophenylacetylene was hydrogenated in isopropanol in the presence of a $\text{Ru}/\text{Al}_2\text{O}_3$ catalyst to obtain 2-methyl-4-(3-aminophenyl)-3-butyne-2-ol. This reference states that is essential to stop the hydrogenation reaction at the stoichiometric point because
25 reduction of the triple bond ensues. In the third step, this aminophenylacetylene was heated in toluene in the presence of sodium hydroxide pellets, with removal of the acetone co-product by distillation, to obtain 3-aminophenylacetylene.

30

This reference also discloses an attempt to similarly react 3-bromoaniline, instead of 3-bromonitrobenzene, with 2-methyl-3-butyne-2-ol to obtain 2-methyl-4-(3-aminophenyl)-3-butyne-2-ol directly, and reports that 3-bromoaniline failed to react at any appreciable rate at temperatures up to ca. 100 °C.

U.S. Patents 4,128,588 and 4,204,078 (each also from the same Sabourin as inventor, with Selwitz) also discloses this preparation of 2-methyl-4-(3-nitrophenyl)-3-butyn-2-ol from 3-bromonitrobenzene and 2-methyl-3-butyn-2-ol. U.S. Patent No. 4,139,561 and *J. Org. Chem.*, vol. 44 (1979), pp. 1223-1236 (both Onopchenko as well as the same Sabourin and Selwitz) also discloses the preparation of 2-methyl-4-(3-aminophenyl)-3-butyn-2-ol from 2-methyl-4-(3-nitrophenyl)-3-butyn-2-ol by hydrogenation with a ruthenium catalyst, and its subsequent conversion to 3-aminophenylacetylene. Both the patent and journal disclosures highlight the difficult challenge of selectively hydrogenating the nitro group in the presence of the acetylene group. Subsequent U.S. Patents 4,215,226; 4,216,341; 4,219,679 and a publication *J. Org. Chem.*, vol. 44 (1979), pp. 3671-3674 from Onopchenko, Sabourin, and Selwitz disclose hydrogenations of 2-methyl-4-(3-nitrophenyl)-3-butyn-2-ol to 2-methyl-4-(3-aminophenyl)-3-butyn-2-ol using other hydrogenation catalysts.

Synlett, 1995, pp. 1115-1116 discloses this preparation of 2-methyl-4-(4-nitrophenyl)-3-butyn-2-ol from 4-bromonitrobenzene and 2-methyl-3-butyn-2-ol in high yield using a catalyst system comprising palladium on carbon, triphenylphosphine, and cuprous iodide in the presence of 2.5 equivalents of potassium carbonate in 1:1 1,2-dimethoxyethane:water at 80 °C. This reference also discloses the preparation of 2-methyl-4-(3-aminophenyl)-3-butyn-2-ol in 78% isolated yield from 3-iodoaniline (in place of 4-bromonitrobenzene) using the same system and the same conditions. The reference does not disclose any attempt to similarly react 3-bromoaniline.

OBJECTS OF THE INVENTION

The object of this invention is to provide an economically preferable, effective and efficient process for the preparation aminophenylacetylenes. A further object of this invention is to provide a process for the preparation of aminophenylacetylenes that avoids the inherent challenge of selectively hydrogenating a nitro group in the presence of an acetylene group. Another object of this invention is to provide a process for the preparation of aminophenylacetylenes from aminophenylhalides. The present invention is directed towards one or more of the above objects. Other objects and advantages will become apparent to persons

skilled in the art and familiar with the background references from a careful reading of this specification.

SUMMARY OF THE INVENTION

5

In its most general terms, the present invention provides a process for preparing aminoarylacetylenes comprising reacting an N-arylmethylidene aminoarylhalide with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt to produce a novel N-arylmethylidene aminoarylacetylene. The N-arylmethylidene aminoarylacetylene may be hydrolyzed to remove the N-arylmethylidene group and provide the aminoarylacetylene. The present invention thereby provides a practical process for preparing an aminophenylacetylene comprising reacting a N-benzylidene aminophenylhalide with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt to produce a novel N-benzylidene aminophenylacetylene, and hydrolyzing the N-benzylidene aminophenylacetylene to the aminophenylacetylene. N-benzylidene aminophenylhalides can be readily prepared from the corresponding benzaldehyde and the corresponding aminophenylhalide by methods known in the art, typically in the presence of an acid catalyst. In certain embodiments of the present invention, the N-benzylidene aminophenylhalide is provided in the reaction by the benzaldehyde, the aminophenylhalide, and an acid catalyst. The invention thereby provides a process for the preparation of aminophenylacetylenes from aminophenylhalides that avoids the inherent challenge of selectively hydrogenating a nitrophenylacetylene to an aminophenylacetylene.

10

15

20

25

30

From experiments using a substoichiometric amount of benzaldehyde relative to the aminophenylhalide, it was surprisingly discovered that the remaining free aminophenylhalide in the mixture with the resulting N-benzylidene aminophenylhalide also reacts, providing a mixture of the aminophenylacetylene and the N-benzylidene aminophenylacetylene. Apparently, the substoichiometric amount of the benzylidene group catalyzes the conversion of aminophenylhalide to the aminophenylacetylene. While not intending to be bound by theory, this can be explained by the benzylidene group being transferred, during the reaction, from the initial product, the N-benzylidene aminophenyl-

acetylene, to the unreacted free aminophenylhalide, converting it to the more reactive N-benzylidene aminophenylhalide.

In one preferred embodiment, the present invention provides a process for the preparation of 3-aminophenylacetylene carbinols from 3-aminophenylhalides comprising reacting a N-benzylidene derivative of the 3-aminophenylhalide with a alpha-hydroxy terminal acetylene in the presence of an amine base and a catalyst system comprising a palladium catalyst comprising a phosphorus ligand and a cuprous halide, and hydrolyzing the resulting novel N-benzylidene 3-aminophenylacetylene carbinol to the 3-aminophenylacetylene carbinol. Optionally, a mixture of the 3-aminophenylhalide and the N-benzylidene 3-aminophenylhalide is reacted to provide a mixture of the 3-amino-phenylacetylene carbinol and the corresponding N-benzylidene aminophenylacetylene carbinol. 3-amino-phenylacetylene carbinols may be converted to 3-aminophenylacetylene by methods known in the art. The invention thereby provides an efficient process for the preparation of 3-aminophenylacetylene from 3-aminophenylhalides.

DETAILED DESCRIPTION OF THE INVENTION


Suitable starting materials and intermediates for the preparation of aminoarylacetylenes by the present invention are aminoarylhalides in general, arylcarboxaldehydes in general, terminal acetylenes in general, N-arylmethylidene aminoarylhalides in general, and N-arylmethylidene aminoarylacetylenes in general. The N-arylmethylidene aminoarylhalide and the N-arylmethylidene aminoarylacetylene may be cis isomers or trans isomers, or mixtures thereof, about the carbon-nitrogen double bond.

Suitable aminoaryl groups for the aminoarylhalide, the N-arylmethylidene aminoarylhalide, the N-arylmethylidene aminoarylacetylene, and the aminoarylacetylene include those in which the aryl ring system is a carbocyclic aromatic ring system, having only carbon atoms in the ring system, and those in which the aromatic ring system is a heterocyclic aromatic ring system, having one or more heteroatoms in the ring system. Typical carbocyclic aromatic ring systems in the aminoaryl group have 6-14 carbon atoms in the aromatic ring system. A preferred carbocyclic aromatic ring systems are phenyl and substituted phenyl groups. Suitable heterocyclic aromatic ring systems in the aminoaryl group have 5-13


atoms in the aromatic ring system which comprises carbon atoms and one or more heteroatoms. Preferred heteroatoms are oxygen, sulfur, and nitrogen. Typical heterocyclic aromatic ring systems have 5 or 6 atoms in an aromatic ring comprising one or more heteroatoms selected from the group oxygen, sulfur, and nitrogen, benz-fused derivatives thereof, and substituted derivatives thereof. Examples of preferred heterocyclic aromatic ring systems in the aminoaryl group include pyridyl, furyl, thiophenyl, pyrrolyl, their benz-fused derivatives quinolinyl, isoquinolinyl, benzfuryl, benzthiophenyl, indolyl, isoindolyl, and substituted derivatives thereof.

Suitable aryl groups for the arylmethylidene group of the arylcarboxaldehyde, the N-arylmethylidene aminoarylhalide, and the N-arylmethylidene aminoarylacetylene include those having the aromatic ring systems described above as suitable for the aminoaryl group. (An arylcarboxaldehyde is an arylmethylidene oxide. For example, benzaldehyde is benzylidene oxide and is phenylcarboxaldehyde and is phenylmethylidene oxide.) The aromatic ring system in the arylmethylidene group may be the same or different from the aromatic ring system in the aminoaryl group.

For the preparation of aminophenylacetylenes via N-benzylidene aminophenylhalides, suitable starting materials and intermediates are aminophenylhalides in general, benzaldehydes in general, terminal acetylenes in general, N-benzylidene aminophenylhalides in general, and N-benzylidene aminophenylacetylenes in general. Suitable aminophenylhalides, benzaldehydes, terminal acetylenes, N-benzylidene aminophenylhalides, and N-benzylidene aminophenylacetylenes include those having the structural formulas I, II, III, IV, and V, respectively. The aminophenylacetylenes prepared from these suitable starting materials and intermediates have the structural formula VI.




I




II




III



IV



V



VI

X in formulas I and IV is a halogen substituent selected from the group consisting of chloro, bromo, and iodo, preferably selected from the group bromo and iodo, and most preferably bromo.

Y in formulas I, IV, V and VI is a substituent selected from substituents that do not interfere with the reaction chemistry of the invention. These are known to persons skilled in the art and can be determined by routine experimentation. Examples of suitable substituents include fluoro, chloro (provided X is bromo or iodo), alkyl (preferably C₁-C₁₂), alkenyl (preferably C₁-C₁₂), alkynyl (preferably C₁-C₁₂), alkoxy (preferably C₁-C₁₂), acyloxy (preferably C₁-C₁₂), aryl, aryloxy, heteroaryl, OH, NO₂, CN, COOH, SO₂R, SOR, NH₂, NH-alkyl (preferably C₁-C₁₂), N-dialkyl (preferably C₁-C₁₂), trihalomethyl, NHCO-alkyl (preferably C₁-C₈), CONH-alkyl (preferably C₁-C₄), CON-dialkyl (preferably C₁-C₄), COO-alkyl (preferably C₁-C₁₂), CONH₂, CO-alkyl (preferably C₁-C₁₂), NHCOH, NHCOO-alkyl (preferably C₁-C₈), CO-aryl, COO-aryl, CHCHCO₂-alkyl (preferably C₁-C₁₂), CHCHCO₂H, PO-diaryl, and PO-dialkyl (preferably C₁-C₈).

The subscript n in the formulas I, II, IV, V, and VI is an integer from 0 to 4, preferably 0 or 1, and most preferably 0. When n=0, no substituent Y is present in the formula. When n is greater than 1, the Y substituents may be the same or different and are selected independently of each other.


Z in formulas II, IV, and V is defined as for Y above. The subscript m is an integer from 0 to 5, preferably 0 or 1, and most preferably 0. When m=0, no substituent Z is present in the

formula. When m is greater than 1, the Z substituents may be the same or different and are selected independently of each other.


The N -benzylidene aminophenylhalides and N -benzylidene aminophenylacetylenes can be trans isomers (as shown in formulas IV and V) or cis isomers, or mixtures thereof, about the carbon-nitrogen double bond.

R in formulas III, V, and VI is hydrogen or any substituent that does not interfere with the reaction chemistry of the invention. These are known to persons skilled in the art and can be determined by routine experimentation. Examples of suitable substituents include alkyl (preferably C_1 - C_{12}), alkenyl (preferably C_1 - C_{12}), alkynyl (preferably C_1 - C_{12}), alkoxy (preferably C_1 - C_{12}), acyloxy (preferably C_1 - C_{12}), aryl, aryloxy, heteroaryl, NH -alkyl (preferably C_1 - C_{12}), N -dialkyl (preferably C_1 - C_{12}), trihalomethyl, $NHCO$ -alkyl (preferably C_1 - C_8), $CONH$ -alkyl (preferably C_1 - C_4), CON -dialkyl (preferably C_1 - C_4), COO -alkyl (preferably C_1 - C_{12}), $CONH_2$, CO -alkyl (preferably C_1 - C_{12}), $NHCOH$, $NHCOO$ -alkyl (preferably C_1 - C_8), CO -aryl, COO -aryl, $CHCHCO_2$ -alkyl (preferably C_1 - C_{12}), and hydroxyalkyl (preferably C_1 - C_{12}).


In a preferred embodiment, a N -benzylidene aminophenylhalide, optionally in mixture with additional aminophenylhalide, is reacted with an α -hydroxy terminal acetylene to provide the N -benzylidene aminophenylacetylene carbinol and, ultimately, the aminophenylacetylene carbinol. R in the α -hydroxy terminal acetylene, the N -benzylidene aminophenylacetylene carbinol and the aminophenylacetylene carbinol is an α -hydroxy-alkyl group of the formula $-C(OH)R'R''$, as shown in formulas VII, VIII, and IX, respectively.



VII



VIII



IX

Y, n, Z, and m in formulas VIII and IX are defined as above.

R' and R" in formulas VII, VIII, and IX can be the same or different and are defined as for
5 R, above. Preferred R' and R" are independently selected from the group consisting of
hydrogen, lower alkyl groups having from 1 to 4 carbon atoms, phenyl, substituted phenyl;
or R' and R" when taken together with the carbon bearing the hydroxyl group form a
saturated cycloalkyl group, preferably a cyclohexyl or cyclopentyl group.

10 The preparation of the alpha-hydroxy terminal acetylenes is well known in the art. For
example, acetylene can be reacted with acetone to form 2-methyl-3-butyn-2-ol (also known
as acetylene dimethylcarbinol), which is a preferred alpha-hydroxy terminal acetylene for
use in the process of this invention. Other suitable alpha-hydroxy terminal acetylenes
include 3-methyl-1-pentyn-3-ol, 3-ethyl-1-pentyn-3-ol, 2-phenyl-3-butyn-2-ol,
15 1-ethynyl-cyclohexanol, and 1-ethynylcyclopentanol.

In the following detailed description of the process, the aminophenylhalide, the
benzaldehyde, the N-benzylidene aminophenylhalide, the N-benzylidene aminophenyl-
acetylene, and the aminophenylacetylene are used, as preferred embodiments, to describe
20 the process for aminoarylhalides in general, arylcarboxaldehydes in general,
N-arylmethylidene aminoarylhalides in general, N-arylmethylidene aminoarylacetylenes in
general, and aminoarylacetylenes in general, respectively.


The N-benzylidene aminophenylhalides can be prepared by methods known in the art for
25 preparing imines from anilines and benzaldehydes, typically in the presence of an acid
catalyst. The N-benzylidene aminophenylhalide can be provided preformed to the reaction
or can be prepared in solution for the reaction from the benzaldehyde, the
aminophenylhalide, and an acid catalyst. The nature and amount of the acid catalyst is not
critical provided it is effective to produce the N-benzylidene aminophenylhalides and does
30 not interfere with the process to form the aminophenylacetylene, which can be determined
by routine experimentation. Benzaldehydes from commercial sources often contain the
corresponding benzoic acid as an impurity, and in many cases, this benzoic acid is all the
acid that is needed to catalyze the formation of the N-benzylidene aminophenylhalides.

Other carboxylic acids (e.g. acetic acid) also provide effective acid catalysts. Hydrohalic acids and amine hydrohalides may also be used. Solid acids, acid resins for example, may also be used.

5 When formed in the solution for reaction, the mole ratio of benzaldehyde to the aminophenylhalide is typically in the range 0.01:1 to 1.1:1, preferably in the range 0.1:1 and 1.0:1. When a mole ratio of 1.0:1 or greater is used, essentially all of the aminophenylhalide is converted to the N-benzylidene aminophenylhalide. When a mole ratio of less than 1.0:1 is used, a mixture of N-benzylidene aminophenylhalide and free
 10 aminophenylhalide is provided. Mixtures of N-benzylidene aminophenylhalide and free aminophenylhalide can also be provided using preformed N-benzylidene aminophenylhalide. The mole ratio of N-benzylidene aminophenylhalide to free aminophenylhalide in such mixtures, whether made using preformed N-benzylidene aminophenylhalide or made using the benzaldehyde is typically in the range 0.01:1 to 0.99:1, preferably in the range
 15 0.1:1 to 0.9:1. As used herein, the term "aminophenylhalide reactant" refers to the N-benzylidene aminophenylhalide in combination with any free aminophenylhalide in mixture with it.

20 The ratio of the aminophenylhalide reactant to the terminal acetylene is not critical. Either reactant may be the limiting reactant and this choice can respond to other considerations, such as which is the more costly reactant to provide and which is more readily separated or removed to an acceptable level from the product. Generally the mole ratio of aminophenylhalide reactant to terminal acetylene is in the range 0.5:1 to 2:1. In typical
 25 embodiments, this ratio is in the range 1:1 to 1.5:1.

The reaction of the aminophenylhalide reactant with the terminal acetylene occurs in the presence of a base according to the following general reaction equation:



30 wherein Ar is the aminophenyl of the aminophenylhalide reactant, and X and R are as defined above. The identity of the base is not critical provided it is effective in the reaction and does not interfere with the reaction, which can be determined by routine

experimentation. Suitable bases include amine bases, carboxylate salts, carbonate salts, and bicarbonate salts. Secondary and tertiary amine bases are preferred. Particularly preferred are dialkyl and trialkyl amines having the general formula $NR^aR^bR^c$, wherein R^a , R^b , and R^c are independently selected from the group consisting of hydrogen and lower alkyl groups having from 1 to 4 carbons; or where two of R^a , R^b , and R^c when taken together with the nitrogen form a heterocyclic amine; with the proviso that no more than one of R^a , R^b , and R^c is hydrogen. Illustrative examples of such preferred amine bases include dimethylamine, trimethylamine, diethylamine, triethylamine, dibutylamine, tributylamine, and N-methylpiperidine. Most preferred are the trialkyl amines. Related amine bases that do not fit this formula precisely, such as N-methylmorpholine, N,N-dimethylaniline, and N,N-dimethylaminopyridine are also suitable. A combination of a soluble base (e.g. an amine base) and an insoluble base (e.g. a carbonate salt) may also be used.

The mole ratio of the base to the limiting reactant, whether the aminophenylhalide reactant or the terminal acetylene is typically at least 1.0. Higher ratios of base to the reactants are suitable. The preferred amine bases may be conveniently used as solvent for the reaction.

Suitable palladium catalysts include those provided by palladium compounds and salts, in particular palladium(0) compounds and palladium(II) compounds and salts. Preferably, the catalyst also comprises a ligand. Suitable ligands include monodentate and bidentate ligands comprising nitrogen or phosphorus as ligating atom. Preferred ligands are triorganophosphine, triorganophosphite, and aromatic nitrogen heterocycle ligands. Examples of preferred ligands include triarylphosphines (e.g. triphenylphosphine), bidentate bis(diarylphosphino) compounds (e.g. 1,1'-bis(diphenylphosphino)butane), trialkylphosphites (e.g. triisopropylphosphite), and pyridine-type ligands (e.g. pyridine, bipyridine). Particularly preferred ligands are triorganophosphines having the general formula $PR^dR^eR^f$, wherein R^d , R^e , and R^f are independently selected from the group consisting of alkyl groups having from 1 to 6 carbon atoms, phenyl, and substituted phenyl groups. The substituents on the phenyl groups can include alkyl groups having from 1 to 4 carbon atoms, alkoxy groups having 1 to 4 carbon atoms, and halogen. Triphenylphosphine is a frequently used preferred ligand.

Suitable and optimal ratios of the ligand to palladium depend on a number of other parameters, including the identity of the ligand, the concentration of the palladium catalyst, the reaction temperature, the reactivity of the reactants, the solvent, and the like, and can be readily determined by routine experimentation. Typically the mole ratio of the ligand to palladium is in the range of 1:1 to 100:1, preferably in the range 2:1 to 50:1, and most preferably in the range 4:1 to 20:1. In certain embodiments, improved conversion of the aminophenylhalide reactant and improved yield of the aminophenylacetylene is obtained by providing an amount of ligand in the reaction mixture in excess of the maximum mole ratio (4:1) that can be bound to the palladium.

The active palladium catalyst may be prepared in advance of its introduction to the reaction mixture, or may be generated in the reaction mixture. It is believed that the active catalyst in the reaction is a palladium(0) compound. The active palladium catalyst may be provided by a preformed ligated palladium(0) compound (e.g. tetrakis(triphenylphosphine)palladium(0)), or may be provided by combining in solution, either *ex situ* or *in situ* to the reaction mixture, a suitable ligand with a suitable palladium(0) compound (e.g. tris(dibenzylideneacetone)palladium(0)). When the catalyst is provided by a palladium(II) compound or salt, the active catalyst is provided by its reduction either *ex situ* or *in situ* to the reaction mixture. Generally, the other components of the reaction mixture (e.g. an amine base) is capable of reducing the palladium(II) to generate the active catalyst *in situ*. This can be determined by routine experimentation. Suitable reductants for *ex situ* generation of the active catalyst from and palladium(II) sources are known in the art and include organomagnesium halide reagents (e.g. methylmagnesium halide) and various hydride reagents (e.g. sodium bis(2-methoxyethoxy)aluminum dihydride). Preferably the palladium(II) is combined with ligand prior to its reduction. The palladium(II) may be provided as a preformed ligated palladium(II) compound (e.g. dichlorobis(triphenylphosphine)palladium(II)) or may be provided by combining in solution a suitable ligand with a suitable palladium(II) compound (e.g. dichlorobis(acetonitrile)palladium(II)) or palladium(II) salt. Suitable palladium(II)-salts include the salts having the general formula PdA_2 , wherein A is an inorganic or organic salt anion. The identity of the anion A is not critical but it must not interfere with the reaction, which can be determined by routine experimentation. Preferred palladium(II) salts include the chlorides, bromides,

carboxylates (e.g. formate, acetate, stearate) and acetylacetonates. Particularly preferred are palladium diacetate and palladium dichloride.

The amount of palladium catalyst is not critical, but should be a catalytic mole ratio less than about 1:10 to the aminophenylhalide reactant, and preferably less than about 1:100. The minimum amount of palladium catalyst depends on other parameters, including the identity of the ligand, the concentration of the ligand, the reaction temperature, the reactivity of the specific reactants, how much of the aminophenylhalide reactant is the N-benzylidene aminophenylhalide, the concentration of the reactants, the solvent, and the maximum time allowed for completion of the reaction, and can be readily determined by routine experimentation. In typical embodiments, a suitable mole ratio of the palladium catalyst to the aminophenylhalide reactant is in the range of 1:10,000 to 1:100, preferably in the range 1:5000 to 1:500.

Suitable cuprous salts include those having the general formula CuA , wherein A is defined as above, but is independently selected. Preferred cuprous salts are the cuprous halides. Cuprous iodide is particularly preferred. The amount of cuprous salt is not critical, but should be a catalytic mole ratio less than about 1:10 to the aminophenylhalide reactant. The minimum amount of cuprous salt depends on other parameters, like those listed above for the amount of palladium catalyst. In typical embodiments, a suitable mole ratio of cuprous salt to palladium catalyst is in the range of 1:1 to 100:1, preferably in the range of 5:1 to 25:1.

The reaction of the aminophenylhalide reactant with the terminal acetylene may be conducted without solvent, with an excess of amine base as solvent, with an additional solvent that is reaction-inert, or with a mixture of excess amine base and a solvent that is reaction inert. By reaction-inert solvent is meant a solvent system which does not react with the reactants or products of the reaction, or react unfavorably with the catalyst. The term solvent system is used to indicate that a single solvent or a mixture of two or more solvents can be used. Representative solvents are aromatic hydrocarbons such as benzene, toluene, xylene; aliphatic hydrocarbons such as pentane, hexane, heptane; acetonitrile; dialkyl ethers; cyclic ethers, polar aprotic solvents such as dimethylformamide, dimethylacetamide, N-methylpyrrolidone, and sulfolane, chlorinated hydrocarbons such as

methylen chloride, dichloroethylene, carbon tetrachloride, and chloroform, and mixtures thereof. The solvent system used need not bring about complete solution of the reactants. Preferred solvents include the amine base and mixtures of the amine base and a hydrocarbon solvent.

5

The reaction temperature is not critical, but is preferably sufficient for the reaction to proceed at a practical rate. Suitable and optimal reaction temperatures depend on a number of other parameters, including the reactivity of the specific catalyst system, the concentration of the catalyst components, the concentrations and reactivities of the specific reactants, and the solvent, and can be readily determined by routine experimentation. In typical embodiments, the reaction is conducted at a temperature in the range from about 20°C to 200°C, preferably from about 50°C to 120°C. It is often convenient to conduct the reaction at the reflux temperature of the reaction mixture.

10

15

The order of addition of the reaction components is not critical. All the reaction components can be added prior to any heating to the reaction temperature, or one or more components may be added when the other components have been brought to the desired reaction temperature. The preferred order of addition for any specific embodiment can be determined by routine experimentation with a view towards both reaction performance and chemical engineering considerations.

20

N-benzylidene aminophenylacetylene formed by the reaction may be hydrolyzed to the aminophenylacetylene by methods known in the art for the hydrolysis of C,N-diaryl imines in general. Typically this is accomplished by treating the imine with water and an acid.

25

The hydrolysis of the N-benzylidene aminophenylacetylene can be conducted at any point in the process subsequent to the conversion of the aminophenylhalide reactant being judged suitably complete. Water and acid can be added to the converted reaction mixture to effect the hydrolysis prior to any separations. Alternatively, the hydrolysis can be conducted later in the separations scheme. Another alternative is to isolate the N-benzylidene aminophenylacetylene, optionally in combination with any free aminophenylacetylene co-product of the reaction, and subsequently subject it to hydrolysis to the aminophenylacetylene.

30

The N-benzylidene aminophenylacetylene and the aminophenylacetylene can each be recovered and isolated by known methods.

EXAMPLES OF THE INVENTION

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following specific examples are intended merely to illustrate the invention and not to limit the scope of the disclosure or the scope of the claims in any way whatsoever.

Example 1

Preparation of 3-(N-benzylidene)aminophenylacetylene dimethylcarbinol:

A solution of equimolar amounts of 3-bromoaniline (1.72 g, 10 mmol) and benzaldehyde (1.06 g, 10 mmol), and benzoic acid (0.20 g, 1.6 mmol) in triethylamine (5 g) was heated at reflux for 12 hours. The reaction was essentially complete by thin-layer chromatography (40 % ethyl acetate/heptane; silica). The ¹H-NMR spectrum of an aliquot showed the presence of N-benzylidene-3-aminophenylbromide (benzylidenyl proton singlet at δ 8.43 (CD₂Cl₂); compared to authentic N-benzylidene aniline at 8.45).

The reaction mixture was cooled and PdCl₂ (2 mg, 0.01 mmol) and triphenylphosphine (73 mg, 0.28 mmol) were added. The mixture was deaerated by bubbling nitrogen. This reaction mixture was heated at reflux for 20 min and then treated with a slurry of CuI (20 mg, 0.1 mmol) in 2-methyl-3-butyn-2-ol (1.1 g, 13 mmol). The reaction mixture was heated at reflux for 12 hours. Thin-layer chromatography (40 % ethyl acetate/heptane; silica) showed the reaction to be complete.

An aliquot of the reaction mixture was dissolved in CH₂Cl₂/hexane and extracted with water. The organic phase was evaporated under a stream of nitrogen. The ¹H-NMR spectrum of the resulting oil was consistent with the benzylidene imine of 3-methyl-4-(3-aminophenyl)-3-butyn-2-ol (N-benzylidene 3-aminophenylacetylene dimethylcarbinol), having resonances for both the benzylidenyl proton (δ 8.46, s, 1H) and the dimethylcarbinol moiety (δ 1.60, s, 6H). ¹H-NMR (CD₂Cl₂): δ 1.60 (s, 6H), 2.5 (brs,

1H), 7.15-7.20 (m, 1H), 7.25-7.49 (m, 3H), 7.45-7.55 (m, 3H), 7.88-7.95 (m, 2H), 8.46 (s, 1H).

Another aliquot of the reaction mixture was partitioned between 40% ethyl acetate/heptane and water. The organic phase was analyzed by GC/MS, showing only one major non-solvent peak. The mass spectrum of this elute showed a molecular ion (M^+) of mass 263, consistent with $C_{18}H_{17}NO$ for N-benzylidene 3-aminophenylacetylene dimethylcarbinol.

Another aliquot of the reaction mixture was hydrolyzed to remove the N-benzylidene group by partitioning between 40 % ethyl acetate/heptane and aqueous HCl. The acidic aqueous phase was separated, basified with aqueous sodium hydroxide, and extracted with 40% ethyl acetate/heptane. Thin-layer chromatography of the organic solution showed (vs. authentic standard) 3-methyl-4-(3-aminophenyl)-3-buten-2-ol as the major component.

This example illustrates the process of the present invention wherein a N-benzylidene aminophenylacetylene (N-benzylidene 3-aminophenylacetylene dimethylcarbinol) is prepared by reacting a N-benzylidene aminophenylhalide (N-benzylidene 3-aminophenylbromide), preformed *ex situ*, with a terminal acetylene (2-methyl-3-buten-2-ol; also called acetylene dimethylcarbinol) in the presence of a base (triethylamine) and a catalyst system comprising a palladium catalyst and a cuprous salt.

Example 2

Preparation of 3-aminophenylacetylene dimethylcarbinol:

To a nitrogen purged 250 mL three-neck flask equipped with a mechanical stirrer, reflux condenser and septum inlet was added benzoic acid (300 mg, 2.8 mmol) and $PdCl_2$ (25 mg). Then 3-bromoaniline (25 g, 145 mmol), benzaldehyde (4.6 g, 44 mmol) and triethylamine (80 mL) were charged. The resulting mixture was vacuum deaerated (150 torr; N_2 backfill; 5x) and heated under nitrogen at reflux for one hour to form the N-benzylidene 3-aminophenylbromide.

The resulting mixture was cooled to 50 °C and triphenylphosphine (0.94 g in 10 mL triethylamine) was added via syringe. Then 2-methyl-3-butyn-2-ol (16 g) was added followed by CuI (0.3 g in 20 mL triethylamine). The reaction mixture was refluxed for 10 hours. After cooling, toluene (20 mL) and water (40 mL) containing 50% sodium hydroxide (11.6 g) were added. The aqueous phase was separated and the resulting mixture was distilled under vacuum (200 torr) while adding more toluene (3x, 75 mL). A total of 300 mL of distillate was obtained. The resulting slurry was cooled and treated with HCl (100 mL, containing 21 g of 37% HCl). This mixture was stirred for 1 hour at room temperature to hydrolyze the N-benzylidene 3-aminophenylacetylene dimethylcarbinol.

The aqueous solution of the product was separated and treated with toluene (35 mL) and ethyl acetate (30 mL). This mixture was cooled (ice bath) and basified with 50% sodium hydroxide (17 g). Some of the aqueous phase was separated and then the mixture was heated to 60 °C. The remaining aqueous phase was separated and more toluene (155 mL) was added. This mixture was cooled to crystallize the product. Filtration and washing with cold toluene and drying *in vacuo* afforded 19.21 g (75.5%) of 2-methyl-4-(3-aminophenyl)-3-butyn-2-ol. ¹H-NMR (DMSO-*d*₆): δ 1.44 (s, 6H), 5.14 (brs, 2H), 5.37 (s, 1H), 6.5-6.7 (m, 3H), 6.97 (t, 1H, J=7.8Hz).

This example illustrates the process of the invention wherein a mixture of an aminophenylhalide and a corresponding N-benzylidene aminophenylhalide, preformed *ex situ* by reacting the aminophenylhalide with a substoichiometric amount of a benzaldehyde (0.3 eq in this case), are reacted with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt to provide a mixture of the aminophenylacetylene and the corresponding N-benzylidene aminophenylacetylene, and hydrolyzing the N-benzylidene aminophenylacetylene in the mixture to the aminophenylacetylene.

Comparative Example

The procedure was the same as in Example 2 using benzaldehyde free of benzoic acid, through the steps intended for the *ex situ* formation of the N-benzylidene 3-aminophenylbromide and the catalyzed reaction with 2-methyl-3-butyn-2-ol, with the

exception that no benzoic acid was added. Upon the addition of the CuI, an apparent precipitation of Pd was observed (not seen in the procedure of Example 2). After several hours of reflux no reaction to form an aminophenylacetylene dimethylcarbinol, with or without the N-benzylidene moiety, could be observed by thin-layer chromatography.

5

Comparison with Examples 2 demonstrates that when the N-benzylidene imine of the aminophenylhalide is not present, in this case because no acid catalyst was provided for its formation from the benzaldehyde, the catalytic reaction to couple the aminophenylhalide with the terminal acetylene does not readily proceed. It further indicates that, in the present invention, the availability of the N-benzylidene aminophenylhalide for reaction serves to stabilize the palladium catalyst.

10

Example 3

Preparation of 3-aminophenylacetylene dimethylcarbinol:

15

A mixture of 4-bromoaniline (1.72 g, 10 mmol), benzaldehyde (0.27 g free of benzoic acid, 2.5 mmol), benzoic acid (20 mg, 0.16 mmol), PdCl₂ (2 mg) and triethylamine (3.5 g) was deaerated by bubbling nitrogen and then heated to reflux for 30 min. The reaction mixture was cooled to 50° C and treated with triphenylphosphine (73 mg in 0.5 g of TEA) followed by 2-methyl-3-butyne-2-ol (1.1 g, 13 mmol) and then CuI (20 mg in 1 g TEA). This mixture was heated at reflux for 12 hr. Thin layer chromatography showed the reaction to form 3-aminophenylacetylene dimethylcarbinol, in mixture with N-benzylidene 3-aminophenylacetylene dimethylcarbinol, to be complete.

20

25

Comparison to the Comparative Example above demonstrates that the addition of an acid catalyst (in this case benzoic acid) in order to form the N-benzylidene aminophenylhalide when using benzaldehyde free of benzoic acid provides for a successful conversion of the aminophenylhalide, in mixture with the N-benzylidene aminophenylhalide, to the aminophenylacetylene, in mixture with the N-benzylidene 3-aminophenylacetylene.

30

Example 4Preparation of 3-aminophenylacetylene dimethylcarbinol:

A mixture of 3-bromoaniline (1.72 g, 10 mmol), 2-methyl-3-butyn-2-ol (1.1 g, 13 mmol), benzaldehyde (0.3 g, 2.8 mmol, containing some benzoic acid impurity), triethylamine (3 g), toluene (3 g), dichlorobis(triphenylphosphine)palladium (15 mg), and triphenylphosphine (80 mg) was deaerated by bubbling nitrogen and heated to 80 °C and treated with CuI (10 mg). This mixture was refluxed for 7 hours. The cooled reaction mixture was treated with aq NaOH (3 mL, 5 mM). The aqueous phase was separated and the organic phase was distilled to remove TEA with more toluene being added. The resulting toluene solution was treated with aq HCl (5 mL, 9%). After 30 min the aqueous phase was separated and basified with 50% NaOH. The precipitated solid was filtered and dried to afford 1.2 g of 3-aminophenylacetylene dimethylcarbinol.

This example demonstrates that the N-benzylidene aminophenylhalide need not be preformed *ex situ*, but can be formed *in situ* from the aminophenylhalide and the benzaldehyde, in this case substoichiometric benzaldehyde containing benzoic acid impurity.

Example 5Preparation of 3-aminophenylacetylene dimethylcarbinol:

A mixture of 3-bromoaniline (1.72 g, 10 mmol), 2-methyl-3-butyn-2-ol (1.1 g, 13 mmol), 3-nitrobenzaldehyde (0.5 g, 3.3 mmol), triethylamine (5 g), dichlorobis(triphenylphosphine)palladium (15 mg) and triphenylphosphine (80 mg) was deaerated by bubbling nitrogen. The reaction mixture was heated to 80 °C and treated with CuI (10 mg). After 7 hr at reflux thin-layer chromatography showed nearly complete conversion of the 3-bromoaniline to a mixture of 3-aminophenylacetylene dimethylcarbinol and N-(3-nitro-benzylidene) aminophenyl acetylene dimethylcarbinol.

This example demonstrates the use of another benzylidene group in the process of the invention, in this case 3-nitrobenzylidene provided by a substoichiometric amount of

3-nitrobenzaldehyde with *in situ* formation of the N-(3-nitrobenzylidene) aminophenylhalide.

Example 6

5 Preparation of 3-aminodiphenylacetylene:

A mixture of 3-bromoaniline (1.72 g, 10 mmol), benzaldehyde (0.27 g, 2.5 mmol, containing some benzoic acid), and PdCl₂ (2 mg) in triethylamine (4 g) was deaerated by bubbling nitrogen. This mixture was refluxed for 45 min, then cooled to 50 °C and treated with triphenylphosphine (73 mg in 0.8 g triethylamine). Phenylacetylene (1.33 g, 13 mmol) and CuI (25 mg in 0.8 g triethylamine) were then added. After refluxing for 6 hr, the reaction mixture was cooled and treated with water (5 mL). The triethylamine layer was concentrated in vacuo and replaced by toluene. To this solution was added aqueous HCl (15 mL, 7 %). A precipitate was formed and it was filtered and washed with toluene. 15 Drying in vacuo afforded 3-aminodiphenylacetylene hydrochloride (1.4 g).

This example demonstrates another terminal acetylene in the process of the invention. Phenylacetylene is used to make an aminodiphenylacetylene.

20 Example 7

Preparation of 5-(4-aminophenyl)-4-pentyn-1-ol:

A mixture of 4-bromoaniline (1.72 g, 10 mmol), benzaldehyde (0.27 g, 2.5 mmol), benzoic acid (0.05 g, 0.4 mmol), PdCl₂ (2 mg) and triethylamine (3.5 g) was deaerated by bubbling 25 nitrogen and then heated to reflux for 45 min. Thin-layer chromatography showed the formation of N-benzylidene 4-aminophenylbromide). The reaction mixture was cooled to 50 °C and treated with triphenylphosphine (73 mg in 1 g of triethylamine) followed by a slurry of CuI (20 mg) and 4-pentyn-1-ol (1.1 g, 13 mmol). This reaction mixture was heated at reflux for 10 hr. The mixture was cooled and diluted with triethylamine (5 mL) 30 and treated with water (6 mL). The aqueous layer was separated and the triethylamine layer was concentrated under vacuum. Toluene was added and the mixture again concentrated. Then aqueous HCl (10 mL, 7%) was added. The resulting aqueous solution was cooled, basified with 50% NaOH, and extracted with ethyl acetate. The ethyl acetate extract was

evaporated under vacuum and the residue was flash chromatographed (20% ethyl acetate - dichloromethane, silica) to yield 0.57 g of 5-(4-aminophenyl)-4-pentyn-1-ol as an oil.

This example demonstrates the use of another aminophenylhalide, 4-bromoaniline, and another terminal acetylene, 4-pentyn-1-ol, in the process of the invention.

Example 8

Preparation of 3-aminophenylacetylene dimethylcarbinol:

A mixture of 3-bromoaniline (1.72 g, 10 mmol), 2-methyl-3-butyn-2-ol (1.1 g, 13 mmol) and benzaldehyde (0.3 g, 2.8 mmol, containing some benzoic acid) in triethylamine (5 g) was treated with palladium diacetate (11 mg) and triphenylphosphine (197 mg). This mixture was deaerated by bubbling nitrogen and then heated to reflux. CuI (10 mg) was added and the mixture was heated at reflux for 7 hours. Product isolation analogous to Example 2 gave 1.2 g of 3-aminophenylacetylene dimethylcarbinol.

This Example demonstrates the use of another palladium salt, palladium diacetate, to provide the palladium catalyst in the present invention.

The present invention has been shown by both description and examples. The Examples are only examples and cannot be construed to limit the scope of the invention. One of ordinary skill in the art will envision equivalents to the inventive process described by the following claims which are within the scope and spirit of the claimed invention.

CLAIMS


1. A process for the preparation of an N-arylmethylidene aminoarylacetylene comprising reacting a N-arylmethylidene aminoarylhalide with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt.
5
2. A process for the preparation of an aminoarylacetylene wherein the N-arylmethylidene aminoarylacetylene produced by the process of claim 1 is hydrolyzed to the aminoarylacetylene.
10
3. The process of claim 1 wherein the N-arylmethylidene aminoarylhalide is provided in the reaction mixture by reacting the corresponding arylcarboxaldehyde and the corresponding aminoarylhalide in the presence of an acid catalyst.
15
4. The process of claim 3 wherein the acid catalyst is an arylcarboxylic acid corresponding to the arylcarboxaldehyde present as an impurity in the arylcarboxaldehyde.
- 20 5. The process of claim 1 wherein the N-arylmethylidene aminoarylhalide is a N-benzylidene aminophenylhalide and the N-arylmethylidene aminoarylacetylene is a N-benzylidene aminophenylacetylene.
- 25 6. A process for the preparation of a aminophenylacetylene wherein the N-benzylidene aminophenylacetylene produced by the process of claim 5 is hydrolyzed to the aminophenylacetylene.
7. The process of claim 5 wherein the terminal acetylene is an alpha-hydroxy terminal acetylene and the N-benzylidene aminophenylacetylene is a N-benzylidene aminophenylacetylene carbinol.
30

8. The process of claim 7 wherein the alpha-hydroxy terminal acetylene is acetylene dimethylcarbinol and the N-benzylidene aminophenylacetylene carbinol is N-benzylidene aminophenylacetylene dimethylcarbinol.
- 5 9. The process of claim 8 wherein the N-benzylidene aminophenylhalide is an N-benzylidene 3-aminophenylhalide and the N-benzylidene aminophenylacetylene dimethylcarbinol is a N-benzylidene 3-aminophenylacetylene dimethylcarbinol.
- 10 10. The process of claim 5 wherein the base comprises a trialkyl amine, wherein the palladium catalyst is selected from catalysts provided by palladium(0) compounds, catalysts provided by palladium(II) compounds, and catalysts provided by palladium(II) salts and comprises a ligand wherein the ligating atom is selected from nitrogen and phosphorus, and wherein the cuprous salt is a cuprous halide.
- 15 11. A process for the preparation of an aminoarylacetylene comprising reacting a mixture of an aminoarylhalide and a corresponding N-arylmethylidene aminoarylhalide with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt to provide a mixture of the aminoarylacetylene and the corresponding N-arylmethylidene aminoarylacetylene.
- 20 12. The process of claim 11 further comprising hydrolyzing the N-arylmethylidene aminoarylacetylene in the mixture to the aminoarylacetylene.
- 25 13. The process of claim 11 wherein the N-arylmethylidene aminoarylhalide is provided in the reaction mixture reacting a portion of the aminoarylhalide provided to the reaction with a substoichiometric amount of the corresponding arylcarboxaldehyde in the presence of an acid catalyst.
- 30 14. The process of claim 13 wherein the acid catalyst is an arylcarboxylic acid corresponding to the arylcarboxaldehyde present as an impurity in the arylcarboxaldehyde.

15. The process of claim 11 wherein the aminoarylhalide is an aminophenylhalide, the corresponding N-arylmethylidene aminoarylhalide is a N-benzylidene aminophenylhalide, the aminoarylacetylene is an aminophenylacetylene, and the corresponding N-arylmethylidene aminoarylacetylene is a N-benzylidene aminophenylacetylene.
16. The process of claim 15 further comprising hydrolyzing the N-benzylidene aminophenylacetylene in the mixture to the aminophenylacetylene.
17. The process of claim 15 wherein the terminal acetylene is an alpha-hydroxy terminal acetylene and the aminophenylacetylene is an aminophenylacetylene carbinol.
18. The process of claim 17 wherein the alpha-hydroxy terminal acetylene is acetylene dimethylcarbinol and the aminophenylacetylene carbinol is an aminophenylacetylene dimethylcarbinol.
19. The process of claim 18 wherein the aminophenylhalide is a 3-aminophenylhalide and the aminophenylacetylene dimethylcarbinol is 3-aminophenylacetylene dimethylcarbinol.
20. The process of claim 15 wherein the base comprises a trialkyl amine, wherein the palladium catalyst is selected from catalysts provided by palladium(0) compounds, catalysts provided by palladium(II) compounds, and catalysts provided by palladium(II) salts and comprises a ligand wherein the ligating atom is selected from phosphorus and nitrogen, and wherein the cuprous salt is a cuprous halide.

21. A process for the preparation of an aminophenylacetylene comprising reacting an aminophenylhalide with a benzaldehyde to form a N-benzylidene aminophenylhalide, reacting the N-benzylidene aminophenylhalide with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt to form a N-benzylidene aminophenylacetylene, and subsequently hydrolyzing the N-benzylidene aminophenylacetylene to form the aminophenylacetylene.
22. A process for the preparation of an aminophenylacetylene comprising reacting an aminophenylhalide with a substoichiometric amount of a benzaldehyde to convert a corresponding amount of the aminophenylhalide to the corresponding N-benzylidene aminophenylhalide, reacting the mixture of the N-benzylidene aminophenylhalide and the remaining aminophenylhalide with a terminal acetylene in the presence of a base and a catalyst system comprising a palladium catalyst and a cuprous salt to form a product mixture of the aminophenylacetylene and the corresponding N-benzylidene aminophenylacetylene, and subsequently hydrolyzing the N-benzylidene aminophenylacetylene in the product mixture to the aminophenylacetylene.
23. An N-arylmethylidene aminoarylacetylene selected from cis isomers and trans isomers, and mixtures thereof, about the carbon nitrogen double bond.
24. The compound of claim 23 wherein the N-arylmethylidene aminoarylacetylene is a N-benzylidene aminophenylacetylene.

25. The N-benzylidene aminophenylacetylene of claim 24 wherein the trans isomer has the formula:



wherein

Y and Z are substituents independently selected from fluoro, chloro, alkyl, alkenyl, alkynyl, alkoxy, acyloxy, aryl, aryloxy, heteroaryl, OH, NO₂, CN, COOH, SO₂R, SOR, NH₂, NH-alkyl, N-dialkyl, trihalomethyl, NHCO-alkyl, CONH-alkyl, CON-dialkyl, COO-alkyl, CONH₂, CO-alkyl, NHCOH, NHCOO-alkyl, CO-aryl, COO-aryl, CHCHCO₂-alkyl, CHCHCO₂H, PO-diaryl, and PO-dialkyl;

n is an integer from 0 to 4, wherein when n is greater than 1, the Y substituents may be the same or different and are selected independently of each other;

m is an integer from 0 to 5, wherein when m is greater than 1, the Z substituents may be the same or different and are selected independently of each other; and

R is selected from hydrogen, alkyl, alkenyl, alkynyl, alkoxy, acyloxy, aryl, aryloxy, heteroaryl, NH-alkyl, N-dialkyl, trihalomethyl, NHCO-alkyl, CONH-alkyl, CON-dialkyl, COO-alkyl, CONH₂, CO-alkyl, NHCOH, NHCOO-alkyl, CO-aryl, COO-aryl, CHCHCO₂-alkyl, and hydroxyalkyl.

26. The compound of claim 25 comprising a N-benzylidene aminophenylacetylene carbinol wherein R is an alpha-hydroxy alkyl group of the formula -C(OH)R'R'' wherein R and R'' are independently selected from hydrogen, lower alkyl groups having from 1 to 4 carbon atoms, phenyl, substituted phenyl; or R' and R'' when taken together with the carbon bearing the hydroxyl group form a cyclohexyl or cyclopentyl group.

27. The compound of claim 26 wherein the N-benzylidene aminophenylacetylene carbinol is N-benzylidene 3-aminophenylacetylene dimethylcarbinol.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/14253

A. CLASSIFICATION OF SUBJECT MATTER IPC(6) :C07C 249/02 US CL :564/272, 273, 274, 275 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 564/272, 273, 274, 275 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched NONE Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAS ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X - Y	BLEICHER et al. Aryl and Heteroaryl-Alkyne Coupling Reactions Catalyzed by Palladium on Carbon and CuI in an aqueous Medium. Synlett. November 1995, Vol. 11, pages 1115-1116, see entire document.	1, 2, 5, 7-11, 15-20 ----- 1, 2, 5, 7-11, 15-20
X - Y	Chem. abstr., Vol. 124, 1996 (Columbus, OH, USA), Whittall et al., 'Organometallic Complexes for Nonlinear Optics. 3. Molecular Quadratic Hyperpolarizabilities of Ene-, Imine-, and Azo-Linked Ruthenium .sigma.Acetylides: X-ray Crystal Structure of Ru(E)-4,4'-C.tplbond.CC6H4CH:CHC6H4NO2)(PPh3)2(.eta.-C5H5), the abstract No. 124:289837, Organometallics. 1996, 15(7), 1935-41, see entire document.	23-25 ----- 1, 2, 9-11, 15, 16, 20
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"B"	earlier document published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	
"P"	document published prior to the international filing date but later than the priority date claimed	"A" document member of the same patent family
Date of the actual completion of the international search 21 DECEMBER 1998		Date of mailing of the international search report 14 JAN 1999
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer <i>Peter G. O'Sullivan</i> PETER G. O'SULLIVAN Telephone No. (703) 308-1235

International application No.
PCT/US98/14253

International application No.
PCT/US98/14253

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X - Y	US 4,730,032 A (ROSSI et al) 08 March 1988, see entire document.	23-25 ----- 1, 2, 9-11, 15, 16, 20
X - Y	EP 0 131 968 A2 (HITACHI, LTD.) 23 January 1985, see entire document.	23-25 ----- 1, 2, 9-11, 15, 16, 20